

KANAZAWA-97-09

**NON-PERTURBATIVE RENORMALIZATION GROUP
APPROACH TO DYNAMICAL CHIRAL SYMMETRY
BREAKING IN GAUGE THEORIES ^a**

Ken-Ichi AOKI

*Department of Physics, Kanazawa University
Kakuma-machi, KANAZAWA 920-11
JAPAN*

We analyze the dynamical chiral symmetry breaking in gauge theories solely by the non-perturbative renormalization group, without recourse to the Schwinger-Dyson method. First, we briefly review the basic notions and formulation, and clarify its great feature that it gives a systematic approximation scheme without any divergent series nor serious gauge dependence, compared to other perturbative and non-perturbative methods. Then we apply this new method to QED and QCD to find that our lowest level approximation improves the ladder Schwinger-Dyson equation results in a gauge independent way. We get the chiral phase structures and the critical exponents in the fixed gauge coupling analysis, and also calculate the chiral condensates in case of running QCD.

1 Introduction

Our aim is to attack the dynamical chiral symmetry breaking solely by the non-perturbative renormalization group (NPRG). The ladder Schwinger-Dyson (SD) equation has been the best analytic tool to investigate it,^{1,2} but is unsatisfactory due to the strong gauge dependence,³ and difficulty of going beyond the ladder,⁴ while the improved ladder approximation with the running gauge coupling constant gives good results in QCD even quantitatively.^{5,6} However the improved ladder is just a model, having no firm theoretical base, thus there has been no way to further improve it.

As everybody knows, NPRG must best fit the issue. We first define a renormalization group equation for some appropriate sub-theory space, then we get flows, critical surfaces and fixed points, and we can calculate critical exponents or anomalous dimensions etc. However nobody had known a good way of approximation to perform this strategy. Recently we found a good approximation, which is a systematic and consistent method, contains the ladder and even the improved ladder SD results, thus we can go beyond the ladder, can improve the improved ladder, in a gauge independent way. It may be applicable to wide range of models and phenomena including supersymmetry, dynamical gauge symmetry breaking, topological effects, etc.

^aInvited talk at International Workshop on Perspectives of Strong Coupling Gauge Theories (SCGT96), Nagoya, 1996. This talk is based on the work in collaboration with K. Morikawa, K. Shimizu, W. Souma, J.-I. Sumi, M. Taniguchi, H. Terao and M. Tomoyose.

In a battle field map of the renormalization group approaches (Fig.1), NPRG is just in the mid of the field, between the perturbative and the non-perturbative head quarters. It shares good features of both sides. The lattice simulation is in fact the most prominent in the non-perturbative regime, but it suffers serious problems of chiral fermions, and that it has exhausted the computer resources on the earth, although recently improved actions enhanced it even thousand times. It is not feasible to simulate systems with large hierarchies. Its great success in QCD is due to the fact that the confinement transition can be described within a relatively short range of energy scales.




Fig.1




Fig.2

On the other hand, in NPRG, there are no difficulty of chiral fermions nor large hierarchies. The most serious problem in NPRG is that it cannot respect the manifest gauge symmetry due to the introduction of the momentum cutoff. However this is not an essential difficulty, but just a practical issue. One should remember that the renormalization calculation in gauge theories could be done even with the momentum cutoff regularization, as was done for QED before the gauge invariant regularization was invented.

Figure 2 shows a more personal view of the field. We, the Wilsonian people, is surrounded by other old countries. In other words we share all beauties of these countries simultaneously. We are now lonely attacking the

weakest neighbor of the republic of ladder also known as Nagoya, and finally will merge it, we hope.

2 Wilsonian renormalization group

In short, NPRG is a method to evaluate integral by solving a corresponding differential equation,

$$F(t) \equiv \int_0^t f(x)dx \implies \frac{\partial F(t)}{\partial t} = f(t), \quad F(0) = 0,$$

which is nothing but a definition of integral, though. We evaluate a path integral by solving a functional differential equation which is called NPRG equation.^{8,9,10,11} The essential ingredient is the Wilsonian effective action S_{eff} defined by

$$\int \mathcal{D}\phi \exp(-S[\phi]) = \int \mathcal{D}\phi_{>} \mathcal{D}\phi_{<} \exp(-S[\phi_{>}, \phi_{<}]) \equiv \int \mathcal{D}\phi_{<} \exp(-S_{\text{eff}}[\phi_{<}]),$$

$$\exp(-S_{\text{eff}}[\phi_{<}]) = \int \mathcal{D}\phi_{>} \exp(-S[\phi_{>}, \phi_{<}]) .$$

The high frequency modes ($\phi_{>}$) are integrated out, and the effective action for the low frequency modes ($\phi_{<}$) is obtained. This is equivalent to the so-called block spin transformation, and it physically describes the coarse graining or the transition from *micro* to *macro*.

Now, we define the flow of S_{eff} by continuously decreasing the cutoff Λ which is the boundary between the high and low frequency modes. The effective action at Λ , $S_{\text{eff}}[\Lambda]$, is obtained after the high frequency modes ($> \Lambda$) have been integrated out: $Z = \int^\Lambda \mathcal{D}\phi \exp(-S[\phi; \Lambda])$, where the measure $\mathcal{D}\phi$ contains low modes only. We evaluate the derivative of $S_{\text{eff}}[\Lambda]$, $\frac{\partial}{\partial \Lambda} S_{\text{eff}}[\phi; \Lambda]$, where the fictitious time parameter t is introduced to control the cutoff: $\Lambda(t) = \exp(-t)\Lambda$, $\delta \Lambda = \Lambda \delta t$. Suppose we lower the cutoff by $\delta \Lambda$, and evaluate the change of the effective action δS_{eff} , which comes from the shell mode ($[\Lambda - \delta \Lambda, \Lambda]$) path integral. We expand S_{eff} with respect to the shell mode field ϕ_{S} ,


$$\begin{aligned} \exp\{-S_{\text{eff}}[\phi_{<}; \Lambda - \delta \Lambda]\} &= \int \mathcal{D}\phi_{\text{S}} \exp\{-S_{\text{eff}}[\phi_{<} + \phi_{\text{S}}; \Lambda]\} \\ &= \exp(-S_{\text{eff}}[\phi_{<}; \Lambda]) \int \mathcal{D}\phi_{\text{S}} \exp\left\{-\int_p' \frac{\delta S_{\text{eff}}}{\delta \phi_p} \phi_{\text{S}}(p)\right\} \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2} \int_p' \int_q' \frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_q} \phi_S(p) \phi_S(q) + \dots \Big\} \\
& = \exp\{-S_{\text{eff}}[\phi_<; \Lambda]\} \int \mathcal{D}\phi_S \\
& \exp \left\{ - \int_p' \left(\frac{\delta S_{\text{eff}}}{\delta \phi_p} \phi_S(p) + \frac{1}{2} \phi_S(p) \frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_{-p}} \phi_S(-p) \right) + O(\delta \Lambda^2) \right\} ,
\end{aligned}$$

where the momentum integration is carried over the shell momentum denoted by the prime. Only the first order term in $\delta \Lambda$ must be kept to evaluate the derivative. The terms with more than two shell modes do not contribute to the first order, since they must be accompanied by at least two loop integral of the shell momentum, thus leaving $O(\delta \Lambda^2)$. Then the shell mode path integral can be exactly carried out, which is just the Gaussian integral, resulting

$$\delta S_{\text{eff}} = \frac{1}{2} \int_p' dp \left\{ -\text{Tr} \ln \left(\frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_{-p}} \right) + \frac{\delta S_{\text{eff}}}{\delta \phi_p} \left(\frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_{-p}} \right)^{-1} \frac{\delta S_{\text{eff}}}{\delta \phi_{-p}} \right\} .$$

Graphically these contributions are represented by the following diagrams, *ring* and *dumbbell*, and its physical reasoning is clear. The shell mode path integral produces all the quantum corrections due to the shell modes, with external fields of the low modes. Note that the tree and the one loop diagrams complete the derivative, and no external momenta may flow in the loop except for the dumbbell endpoint vertices.



To make NPRG equation physically sensible, we transform variables to be dimensionless, where the dimensional unit is taken to be the cutoff at that ‘time’. This transformation runs the coupling constants, which is called the canonical scaling. Also we introduce the wave function renormalization to normalize the kinetic terms of every fields, which chooses a representative of the equivalent classes in the theory space. Then we finally get

$$\begin{aligned}
\frac{\partial S_{\text{eff}}}{\partial t} & = -\frac{1}{2\delta t} \int_p' \left\{ -\text{Tr} \ln \left(\frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_{-p}} \right) + \frac{\delta S_{\text{eff}}}{\delta \phi_p} \left(\frac{\delta^2 S_{\text{eff}}}{\delta \phi_p \delta \phi_{-p}} \right)^{-1} \frac{\delta S_{\text{eff}}}{\delta \phi_{-p}} \right\} \\
& + dS_{\text{eff}} + \int_p \phi_p \left(\frac{2-d-\eta}{2} - p^\mu \frac{\partial'}{\partial p^\mu} \right) \frac{\delta}{\delta \phi_p} S_{\text{eff}} ,
\end{aligned}$$

where d is the space-time dimension, η is the anomalous dimension. This equation, called the Wegner–Houghton equation,⁹ describes the change of the effective action, that is, of a set of infinite number of coupling constants. Thus the right-hand side should be called a β *functional*, which controls the flows in the infinite dimensional theory space. Important is that it is obtained as an exact and closed form, therefore it is sometimes called the exact renormalization group.

3 Approximation method

Even though we obtained the exact renormalization group, it is impossible to solve such an infinite dimensional functional differential equation. Thus what is more important is if there is a good way of systematic approximation or not. The approximation method for the exact renormalization group is a projection to a sub-theory space. For example, we take a sub-theory space spanned only by the local potential terms in addition to the kinetic terms

$$S_{\text{eff}} = \int d^d x \left\{ V(\phi) + \frac{1}{2}(\partial_\mu \phi)^2 \right\} ,$$

which is regarded as the lowest order of the expansion of the effective action with respect to the number of field derivatives. This is called the local potential approximation (LPA),^{9,16} where the corrections to the kinetic terms are ignored either, thus it gives vanishing anomalous dimensions. The Wegner–Houghton equation in LPA is reduced to read

$$\frac{\partial V(\phi, t)}{\partial t} = \frac{A_d}{2} [\ln(1 + V'')] + d \cdot V + \frac{2-d}{2} \phi V' ,$$


where V' denotes the derivative with respect to the field ϕ , and A_d is a constant of the d -dimensional angular integral. We see the original functional differential equation is simplified to a 2-dimensional (t, ϕ) partial differential equation, which is non-linear, and is to be solved as an initial value problem. Note that LPA still deals with an infinite dimensional theory space spanned by an arbitrary function $V(\phi)$.

It is still not easy to analyze the system precisely. We reduce it to a finite dimensional sub-theory space by expanding the potential in terms of the polynomials in ϕ . For example, take a scalar theory with Z_2 symmetry in 3 space-time dimension, and expand the potential as

$$V(\phi, t) = a_1(t)\rho + a_2(t)\rho^2 + a_3(t)\rho^3 \cdots , \quad \rho \equiv \phi^2/2 .$$


Then the β functions (flow equations for the coefficient functions) are given by

$$\begin{aligned}\dot{a}_1 &= 2a_1 + \frac{3}{2\pi^2} \frac{a_2}{1+a_1}, \\ \dot{a}_2 &= a_2 - \frac{9}{2\pi^2} \frac{a_2^2}{(1+a_1)^2} + \frac{5}{\pi^2} \frac{a_3}{1+a_1}, \\ \dot{a}_3 &= \frac{27}{2\pi^2} \frac{a_2^3}{(1+a_1)^3} - \frac{45}{2\pi^2} \frac{a_2 a_3}{(1+a_1)^2}.\end{aligned}$$



Even in this crude approximation with only 3 operators, we find a non-trivial fixed point realizing the ferromagnetic phase transition with the critical exponent $\nu = 0.586$, where the lattice simulation gives 0.629, such a good coincidence. We increase the number of operators or the dimension of the sub-theory space to see if the results converge. If it does, we understand we get the results for LPA sub-theory space.

The approximation here is a projection to a sub-theory space, giving a projected β functions, which define a projected flow. The projected flow is different in general from the projection of the true flow. The true flow always goes out of the sub-theory space. The projected flow pulls it back on the sub-theory space in each step of transformation. In some exceptional case as the large- N theory, the projected flow coincides with the projection of the true flow if we choose a good coordinate system of the theory space, which we call the perfect coordinate.¹²



Generally we may expect that enlarging the sub-theory space, the physical results converge, or oscillate at worst. This is a natural expectation. Enlarging the sub-theory space is not directly related to any divergent series expansion. It is more like enlarging the total lattice size in simulation, where nobody expects the results to diverge. However, usually we must think out a good series of sub-theory spaces to get fast convergence. We have learned that an ‘environmentally friendly’ coordinate system assures better convergence. For example, expanding the potential, we might define the following two schemes: a fixed scheme (A) and a comoving scheme (B),¹⁵

$$\begin{aligned}\text{A: } V(\rho) &= \sum_{n=1}^{nmax} a_n \rho^n & \text{B: } V(\rho) &= \sum_{n=1}^{nmax} a_n (\rho - \rho_0(t))^n\end{aligned}$$

The comoving scheme is a dynamic coordinate since its origin of expansion $\rho_0(t)$ is taken as a minimum of the potential at time t . Physically it is better since it uses more appropriate propagators at t , and it actually improves convergence drastically as is seen in Fig.3.^{12,21,22}^b

4 Comparison with other non-perturbative methods

We compare our NPRG results with other standard non-perturbative methods. First let us examine the leading critical exponent of d -dimensional scalar theory comparing with the ϵ -expansion.¹² As is seen in Fig.4, the ϵ -expansion gives a divergent series. Of course it is extensively investigated that these divergent series can be Borel resummed to give a convergent value which is also displayed in the figure. Our NPRG analysis gives a globally right behavior in the lowest order approximation LPA.




Fig.3




Fig.4

Next we analyze scalar theories with N -components in 3 space-time dimension, and compare NPRG with the $1/N$ expansion (Fig.5).^{12,13,14} The $1/N$ leading result is just $\nu = 1$, and we show the next to leading ($O(1/N)$) and the next next to leading ($O(1/N^2)$) results. It is clearly seen that the $1/N$ expansion is also a divergent series. For small N theories, the leading result ($\nu = 1$ classical result) is the best, and adding higher orders in $1/N$ makes it

^bAfter the conference, we find that even the comoving frame gives finally diverging results. However, it is at extremely large orders, thus it is after the highly accurate convergence of even 8 digits for ν has been obtained²². We thank T. R. Morris to motivate us to investigate very large orders.

worse. Our NPRG analysis gives a stable and globally correct results also in this comparison, and becomes exact at $N \rightarrow \infty$ limit. We expect that when enlarging the sub-theory space beyond LPA, we get closer to the true values. Some of the beyond LPA results are also plotted in Fig.5 and our expectation proves right at least in the small N region. Note that Borel resummation for $1/N$ expansion is not feasible at all.




Fig.5




Fig.6

What is a crucial difference between NPRG and, for example, $1/N$ expansion? The $1/N$ expansion contains only limited structures of diagrams corresponding to the order of the expansion, while NPRG contains every structured diagrams, exactly for the leading, but not exactly for higher orders in $1/N$, even at the lowest level of LPA. Generally NPRG contains every powers of parameters as ϵ , $1/N$, and coupling constants, and it contains the lowest order exactly. This feature assures that NPRG gives globally correct behavior, and it is expected to converge when enlarging the sub-theory space.

We show another striking results in case of scalar theories in 1 space-time dimension also known as the unharmonic oscillator,¹⁷ whose action is given by

$$S = \int dt \left(\frac{1}{2} \dot{x}^2 \pm \frac{1}{2} x^2 + \lambda x^4 \right) .$$

We evaluate the energy gap between the ground state and the first excited state using the infrared effective potential obtained by NPRG. In case of the single-well potential (Fig.7), our results are almost exact for whole range of the

coupling constant λ , whereas the standard perturbation series shows its divergent nature even at the weak coupling region (although Borel resummation is possible in this case).




Fig.7




Fig.8

In case of the double-well potential, we compare our results with the instanton calculation (Fig.8). At $\lambda \sim 0.08$ the first excited state energy crosses down the central barrier, where the LPA results start deviating from the exact values. At this point the instanton results are already unreliable. The LPA works extremely well except for the deep region where the tunneling rate is very small and the dilute instantons become exact.

To see how NPRG works well for the double-well potential is very important, since it is related to a more general issue of how NPRG can treat essential singularities and topological objects in field theory. The LPA gives a convex effective potential in case of the spontaneous symmetry breaking, which indicates that LPA is able to take account of the global domain wall structures without which the convexity may not appear.¹⁷ However it seems that LPA is not successful to evaluate precisely the action accompanying such topological configurations since it fails to give the correct energy gap in case of low tunneling rate.

5 Non-perturbative RGE vs. Perturbative RGE

Here we recapitulate the features of NPRG, compared to the standard perturbative RGE (PRGE). PRGE defines the β functions which are calculated by the loop expansion, giving the leading log series expansion or the improved perturbation, while NPRG β functional is infinite dimensional, but it is exactly given by tree and one-loop diagrams. Multi-loop effects come out through the

higher dimensional operators. PRGE gives the asymptotic series at best, and actually other non-perturbative methods including the ϵ -expansion and the $1/N$ -expansion do as well, while in NPRG the approximation is defined by a projection to a sub-theory space and systematic enlargement of it, thus no diverging series may appear since no expansion in any parameter is employed.

The standard effective action $\Gamma(\phi)$ obtained by PRGE does not serve the dynamical symmetry breaking due to a composite order parameter. We need to evaluate Cornwall-Jackiw-Tomboulis (CJT) effective potential for the composite operators or equivalently to solve the Schwinger-Dyson (SD) equation.⁷ However, NPRG automatically contains the CJT effective potential through the Wilsonian effective action including higher dimensional multi composite operators, thus the non-perturbative vacuum can be directly analyzed.

NPRG cannot manifest the gauge invariance due to the momentum cutoff. Then theory space must span gauge non-invariant operators as well, and we have to pick up the gauge invariant theory space (which differs from the gauge invariant operator space).^{20,18} This is a serious problem for load of the practical calculation, but it is not an essential difficulty and can be treated anyway.

To summarize, NPRG method is very general, robust and global, and no divergent series appears. Recent development has been done by numerical integration of the NPRG equation in LPA and beyond LPA sub-theory spaces. This is a balanced research between analytic and numeric. The lowest order sub-theory space, LPA, has exhibited the features that it contains one-loop perturbative results, exact in the large N leading, exact in the 1st order of ϵ -expansion, and furthermore contains the improved ladder SD results.

6 Dynamical chiral symmetry breaking

Due to shortage of pages I have, we just briefly describe the basic results of our works.¹⁹ We consider a chiral invariant gauge theory and set up NPRG equations in LPA. We include multi-fermi operators, 4-fermi, 8-fermi, $\dots$, respecting the chiral invariance. The β functions for 4-fermi operators do not depend on 8- or more-fermi operators due to the chiral invariance. Thus our multi-fermi coordinates is a perfect coordinates and the flow of 4-fermi operators can be solved exactly in LPA, coupled with the running of the gauge coupling constant e ($\lambda = 3e^2/4\pi^2$):

$$\begin{aligned}\frac{dG_S}{dt} &= -2G_S + \left(2 - \frac{1}{2}\right) G_S^2 - 4G_S G_V + \lambda G_S + \left(\frac{1}{8} + \frac{1}{24}\right) \lambda^2, \\ \frac{dG_V}{dt} &= -2G_V + \frac{1}{4} G_S^2 - \lambda G_V - \frac{1}{12} \lambda^2.\end{aligned}$$

Figure 9 shows the diagrams contributing to the β functions of the scalar 4-fermi G_S and vector 4-fermi G_V interactions, where the dashed lines represent the gauge boson.




Fig.9




Fig.10

We define a part of the β function, the ladder part, as in Fig.9. If one takes only this part in the following analyses, one obtains exactly the same results as those by the ladder SD equation in the Landau gauge. (This uniqueness of the Landau gauge comes from the fact that the fermion anomalous dimension vanishes in this gauge.) Remember that the ladder SD suffers a strong gauge dependence,³ whose reason is clear here. The ladder picks up just a part of the gauge invariant set of diagrams in the β function, taking the ladder but not the crossed ladder. On the other hand, NPRG β function takes both, no difference for either ladder or crossed ladder, thus it respects the gauge independence of flows and of the physical results.

One readily sees (Fig.10) that at $\lambda = 0$, the scalar 4-fermi flows has an ultraviolet fixed point which is nothing but the Nambu-Jona-Lasinio (NJL) criticality, and an infrared fixed point of the Gaussian one. There appear two phases, strong and weak. Note that the 8-fermi operator has no fixed point without gauge interactions, indicating the triviality of the pure fermi theory. When the gauge coupling is switched on, the Gaussian fixed point moves up, the NJL fixed point moves down, and they finally meet together to pair annihilate at a critical gauge coupling λ_c , above which there is only one phase left. To identify these two phases, we investigate the CJT effective potential for the chiral condensate $\langle \bar{\psi}\psi \rangle$, and find that the weak phase is a symmetric

phase, while the strong phase exhibits the spontaneous symmetry breaking of the chiral symmetry, where the potential takes the flat bottomed convex shape characteristic to the spontaneous symmetry breakdown. This structure is displayed as a phase diagram in the scalar 4-fermi vs. the gauge coupling plane in Fig.11, which should be compared with the old traditional phase diagram.² We may calculate a physical quantity of this system, the anomalous dimension of the mass operator $\psi\psi$ (Fig.12). It is enhanced compared to the ladder SD results. Note that our results are LPA exact, including ‘beyond the ladder’ diagrams, and are gauge independent.




Fig.11




Fig.12

We may run the gauge coupling constant. The U(1) gauge interactions keep the two phase structure. However the fixed points are NJL and the Gaussian only, and the renormalized trajectory is on the trivial line, which is consistent with the recent lattice simulations.²³ Asymptotically free gauge interactions wipe out the symmetric phase, and only the Gaussian fixed point survives. Therefore the whole region of the theory space belongs to the chiral symmetry broken phase. We can evaluate physical quantities, the chiral condensate and the fermion mass. In this running gauge coupling case, we proved that the ladder part β function now gives exactly the same results as those by the improved ladder SD equation in the Landau gauge, *a la* Higashijima,⁵ where the scale parameter of the running gauge coupling constant takes a larger momentum of the fermion legs. This particular definition of the running gauge coupling constant exactly emerges from the NPRG LPA with the ladder part. Now we know what is included and what is missing in the improved ladder SD equations, thus establishing the improved ladder for the first time.

To evaluate the infrared physical quantities, the bare multi-fermi coor-

dinate system is not good, since it uses the massless fermion propagators. We introduce a collective coordinate of a scalar field coupled to the $\bar{\psi}\psi$ operator, which is another example of the environmentally friendly coordinate system. Within the ladder part, we have calculated the chiral condensate and the fermion mass as in Fig.13, which shows good convergence within low dimensional sub-theory spaces, and agrees with the improved ladder SD results.⁶ Taking the non-ladder β components into account, we proceed to evaluate infrared physical quantities beyond the ladder SD results, in a gauge independent way. That will be the first results of gauge independent and including ‘beyond the ladder’, thought it is not an easy task.




Fig.13

To deal with the gauge interactions more seriously, we need to formulate NPRG with a smooth cutoff function.^{10,11} Then the cutoff scheme (the profile of the cutoff function) dependence might cause a problem. However we find that indeed such scheme dependence affects phase criticalities (the critical coupling constants) which are themselves not physical quantities, but the physical quantities like the anomalous dimension of the mass operator has only very small scheme dependence.²⁰

We have clarified that NPRG equipped with the sub-theory space approximation potentially solves the dynamical chiral symmetry breaking in gauge theories, which can go beyond the ladder SD results restoring the gauge independence. This new method of treating non-perturbative QCD will give us a promising tool alternative to the lattice simulations. However we may need some revolutionary new techniques of evaluating the β functional to solve QCD

with larger dimensional sub-theory spaces of the gauge sector keeping the gauge invariance.

Acknowledgments

The author is partially supported by Grant-in Aid for Scientific Research (#08240216 and #08640361) from the Ministry of Education, Science and Culture.

References

1. Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345. T. Maskawa and H. Nakajima, Prog. Theor. Phys. **52** (1974) 1326. R. Fukuda and T. Kugo, Nucl. Phys. **B117** (1976) 250. V. A. Miransky, Nuovo Cim. **90A** (1985) 149.
2. K.-I. Kondo, H. Mino and K. Yamawaki, Phys. Rev. **D39** (1989) 2430. K. Yamawaki, in *Proc. Johns Hopkins Workshop on Current Problems in Particle Theory 12, Baltimore, 1988*, eds. G. Domokos and S. Kovesi-Domokos (World Scientific, Singapore, 1988). T. Appelquist, M. Soldate, T. Takeuchi and L. C. R. Wijewardhana, *ibid.* W. A. Bardeen, C. N. Leung and S. T. Love, Phys. Rev. Lett. **56** (1986) 1230. C. N. Leung, S. T. Love and W. A. Bardeen, Nucl. Phys. **B273** (1986) 649.
3. K.-I. Aoki, M. Bando, T. Kugo, K. Hasebe and H. Nakatani, Prog. Theor. Phys. **81** (1989) 866.
4. K. Kondo and H. Nakatani, Nucl. Phys. **B351** (1991) 236; Prog. Theor. Phys. **88** (1992) 7373. K. Kondo, Int. J. Mod. Phys. **A7** (1992) 7239.
5. K. Higashijima, Phys. Rev. **D29** (1984) 1228.
6. K.-I. Aoki, T. Kugo and M. G. Mitchard, Phys. Lett. **B266** (1991) 467. K.-I. Aoki, M. Bando, T. Kugo, M. G. Mitchard and H. Nakatani, Prog. Theor. Phys. **84** (1990) 683.
7. J. M. Cornwall, R. Jackiw and E. Tomboulis, Phys. Rev. **D10** (1974) 2428.
8. K. G. Wilson, I. G. Kogut, Phys. Rep. **12** (1974) 75.
9. F. Wegner, A. Houghton, Phys. Rev. **A8** (1973) 401.
10. J. Polchinski, Nucl. Phys. **B231** (1984) 269. G. Keller, C. Kopfer and M. Salmhofer, Helv. Phys. Acta **65** (1992) 32.
11. C. Wetterich, Phys. Lett. **B301** (1993) 90. M. Bonini, M. D'Attanasio, G. Marchesini, Nucl. Phys. **B409** (1993) 441. T. R. Morris, Int. J. Mod. Phys. **A9** (1994) 2411.

12. K-I. Aoki, K. Morikawa, W. Souma, J.-I. Sumi and H. Terao, Prog. Theor. Phys. **95** (1996) 409.
13. W. Souma, private communication.
14. T. R. Morris, hep-th/9610012, in *Proc. of RG96, Dubna, Russia*. T. R. Morris and M. D. Turner, hep-th/9704202.
15. M. Alford, Phys. Lett, **B336** (1994) 237. N. Tetradis and C. Wetterich, Nucl. Phys. **B422** (1994) 541. T. R. Morris, Phys. Lett. **B329** (1994) 241.
16. A. Hazenfratz and P. Hazenfratz, Nucl. Phys. **B270** (1986) 269. T. R. Morris, Phys. Lett, **B334** (1994) 355.
17. K-I. Aoki, H. Terao and M. Taniguchi in preparation.
18. C. Becchi, *On the construction of renormalized quantum field theory using renormalization group techniques*, in: Elementary Particles, Field Theory and Statistical Mechanics, eds. M. Bonini, G. Marchesini, and E. Onofri, Parma University, 1993. U. Ellwanger, Phys. Lett. **B335** (1994) 364. U. Ellwanger, M. Hirsch, and A. Weber, Z. Phys. **C69** (1996) 687. M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. **B418** (1994) 81; **B421** (1994) 429; **B437** (1995) 163; Phys. Lett. **B346** (1995) 87. M. D'Attanasio, and T. R. Morris, Phys. Lett. **B378** (1996) 213. F. Freire, and C. Wetterich, Phys. Lett. **B380** (1996) 337.
19. K-I. Aoki, K. Morikawa, K. Shimizu, J.-I. Sumi, H. Terao and M. Tomoyose, KANAZAWA-96-21. K-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, KANAZAWA-96-22.
20. K-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, Prog. Theor. Phys. **97-3** (1997) 479.
21. R. D. Ball, P. E. Haagensen, J. I. Latorre and E. Moreno, Phys. Lett. **B347** (1995) 80.
22. K-I. Aoki, K. Morikawa, W. Souma, J.-I. Sumi and H. Terao, in preparation.
23. Talk in this conference by G. Schierholz.